arXiv:1206.5205v2 [quant-ph] 6 Jul 2012

Quantum Information Processing and Relativistic Quantum Fields

Dionigi M. T. Benincasa[1], Leron Borsten[1], Michel Buck[1], Fay Dowker[1,2,3]
[1]*Blackett Laboratory, Imperial College, London, SW7 2AZ, U.K.*
[2]*Perimeter Institute for Theoretical Physics, 31 Caroline St. N., Waterloo ON, N2L 2Y5, Canada and*
[3]*Institute for Quantum Computing, University of Waterloo, ON, N2L 3G1, Canada*

It is shown that an ideal measurement of a one–particle wave packet state of a relativistic quantum field in Minkowski spacetime enables superluminal signalling. The result holds for a measurement that takes place over an intervention region in spacetime whose extent in time in some frame is longer than the light crossing time of the packet in that frame.

PACS numbers: 03.67.-a, 03.65.Ta, 03.70.+k

I. INTRODUCTION

In a typical quantum information processing scheme classical agents use a quantum system to encode, process and communicate information (classical bits). A demand to be made of a description of a quantum information processing (QIP) protocol, then, is: show me the bits. In other words it must be understood how the classical bits are encoded physically in the quantum system and how they are read out. A qubit is a two-dimensional quantum system which can be a building block of the total quantum system which is used to perform the information processing task. In a typical protocol external agents feed bits into the system by preparing some of the qubits in one quantum state or another and read bits out at the end by making measurements on some (other) of the qubits. Intermediate steps may involve the performance of unitary transformations and other sorts of operations on the qubits by the agents. So a related demand is: show me the physical qubits. This is particularly important in relation to entanglement. After all, given *any* pure state, $|\psi\rangle$, in a 4-dimensional Hilbert space, there is a decomposition of that Hilbert space as a tensor product of two 2-dimensional spaces – two 'qubits' – in which $|\psi\rangle$ is a maximally entangled state as expressed in the product basis. These 'qubits' will generally be useless for QIP because they will not correspond to any physically accessible two-dimensional systems that can be manipulated by external agents. Thus, in assessing the relevance of calculations of measures of entanglement between qubits for QIP, it is necessary to determine whether and how external agents can intervene upon them.

The satisfaction of these demands to identify how bits are encoded into and read out from useful qubits and how those qubits can be operated upon physically becomes particularly challenging when investigating QIP in a relativistic spacetime, taking into account the locations in spacetime of the actions of the external agents on the quantum system. An obvious framework for such an investigation is relativistic quantum field theory. Here, progress is hampered by the lack of a universally applicable rule for calculating the probabilities of the outcomes of ideal measurements on a relativistic quantum field in a collection of spacetime regions. Indeed, a straightforward relativistic generalisation of the non-relativistic formula for these probabilities leads to superluminal signalling [1].

We review this generalised rule. We work in the Heisenberg (Interaction) Picture. Let $\mathcal{O}_i$, $i = 1, \ldots n$ be a collection of regions in a globally hyperbolic spacetime. Let $\preceq$ be a relation on the regions defined by $\mathcal{O}_j \preceq \mathcal{O}_k$ iff some point in $\mathcal{O}_j$ is in the causal past of some point in $\mathcal{O}_k$. The transitive closure of this relation is taken and the resulting relation denoted by the same symbol $\preceq$. If the resulting relation is acyclic – *i.e.* $\mathcal{O}_j \preceq \mathcal{O}_k$ and $\mathcal{O}_k \preceq \mathcal{O}_j$ implies $j = k$ – then it is a partial order and there exists a linear ordering of the regions which is compatible with the partial order. In other words there is an assignment of labels $i = 1, \ldots n$ to the regions such that $\mathcal{O}_j \preceq \mathcal{O}_k$ implies $j \leq k$. We assume from now on that the regions and the labelling satisfy this condition. Consider, for each i, the measurement of an observable A_i in region $\mathcal{O}_i$ and consider a particular possible outcome of that measurement corresponding to a (Heisenberg) projection operator, P_i say, onto the relevant eigenspace of the observable. The region $\mathcal{O}_i$ is the "intervention region" in which the external agents act upon the field to effect the measurement of A_i. Note that the intervention region, $\mathcal{O}_i$ can be larger but not (presumably) smaller than the minimal region in which observable A_i is defined. The probability of obtaining those particular outcomes to the sequence of measurements in the regions $\mathcal{O}_i$ is proposed to be [1]

$$\mathrm{Tr}\,(P_n \ldots P_1 \rho P_1 \ldots P_n) \tag{1}$$

where ρ is the initial state.[1]

II. IMPOSSIBLE MEASUREMENTS OF WAVE PACKETS

Henceforth, we will explicitly consider $(d+1)$-dimensional Minkowski space with a mostly plus metric convention, and a free massless scalar field $\hat{\phi}(x)$, though much of what follows will apply with little or no modification for free massive fields and the general conclusions are relevant in globally hyperbolic spacetimes. Using the formula (1), the following protocol will generally result in superluminal signalling [1]. There are three intervention regions: $\mathcal{O}_1$ is a spacetime point X^μ where $X^0 \leq 0$, $\mathcal{O}_2$ a slab of spacetime between $t = 0$ and $t = T > 0$ and $\mathcal{O}_3$ is spacetime point Y^μ spacelike to X and with $Y^0 \geq T$. The interventions in these regions are, respectively, a unitary transformation $e^{i\lambda\hat{\phi}(X)}$ where λ is a real number, a measurement of $|1\rangle\langle 1|$ where $|1\rangle$ is a one particle state and a measurement of $\hat{\phi}(Y)$. The expected value of the outcome of the measurement of $\hat{\phi}(Y)$ depends on λ and the strength of the signal is quantified by [1]

---

[1] If one wants to interpret the rule as corresponding to an effect on the quantum state, it would be that for each intervention region $\mathcal{O}_i$ in turn – in the linear order defined by the labels, $i = 1, \ldots n$ – the quantum state collapses along the boundary of the causal past of the region $\cup_{k=1}^{i} \mathcal{O}_k$.

$$S(X, Y) := -\mathrm{Im}\left(\psi(X)^*\psi(Y)\right) \tag{2}$$

where $\psi(\xi) := \langle 0|\hat{\phi}(\xi)|1\rangle$ is the "one–particle wave-function." $S(X, Y)$ is the derivative of the expected outcome of the measurement of $\hat{\phi}(Y)$ w.r.t. λ, at $\lambda = 0$. Fixing spacetime point X so that $\psi(X)$ is real and nonzero, we obtain $\mathrm{Im}\left(\psi(Y)\right)$ as a measure of the strength of the superluminal signal.

It is straightforward to show that this can be nonzero when the state $|1\rangle$ is a one particle state with a precise d-momentum, $\mathbf{k}$. (Spatial d-vectors will be written in boldface.) Moreover, in this case the wave-function satisfies $\psi(Y^\mu + \xi^\mu) = \psi(Y^\mu)$ where ξ^μ is any null vector proportional to the 4-momentum, $k^\mu = (|\mathbf{k}|, \mathbf{k})$. So the superluminal signal remains no matter how large T, no matter how long the measurement-intervention takes. We conclude that an ideal measurement of this single momentum one–particle state is impossible, if superluminal signalling is forbidden (and assuming that the interventions at X and Y can be done). Such a result is not surprising, given the nonlocal character of a fixed momentum state: it is defined on an entire spacelike hypersurface. What might be more surprising is that similar conclusions can be drawn for ideal measurements of localised one–particle wave packet states as we will now show.

Consider the Gaussian one–particle state

$$|1_d\rangle := (\pi\sigma^2)^{-\frac{d}{4}} \int d^d k \, e^{-\frac{(\mathbf{k}-\mathbf{k}_0)^2}{2\sigma^2}} a^\dagger_{\mathbf{k}}|0\rangle \tag{3}$$

where $a^\dagger_{\mathbf{k}}$ is the creation operator for a one particle state with d-momentum $\mathbf{k}$, $\mathbf{k}_0$ is the mean momentum, σ is the spread in momentum space and $|\mathbf{k}_0| \gg \sigma$.

Consider first $d = 1$ and let $k_0^\mu = k_0(1, 1)$ where $k_0 > 0$, so the packet is moving in the positive space direction. Then for any null vector $\xi^\mu \propto k_0^\mu$ we have:

$$\begin{aligned}\psi(Y^\mu + \xi^\mu) &= \langle 0|\hat{\phi}(Y^\mu + \xi^\mu)|1_1\rangle \\ &= (\pi\sigma^2)^{-\frac{1}{4}} \int_{-\infty}^{\infty} \frac{dk}{4\pi|k|} e^{-\frac{(k-k_0)^2}{2\sigma^2}} e^{ik_\mu(Y^\mu+\xi^\mu)} .\end{aligned} \tag{4}$$

To avoid the pole at the origin, let us modify the wavepacket slightly, allowing it to have support in momentum space only for $k > \epsilon > 0$ where ϵ is small. Then for all momenta contributing to the integral we have $k_\mu\xi^\mu = 0$ and, hence,

$$\psi(Y^\mu + \xi^\mu) = \psi(Y^\mu) . \tag{5}$$

In $1 + 1$ dimensions, when the packet has support only on momenta in the positive spatial direction, it holds its form, does not spread and the superluminal signal persists undiminished for arbitrary intervention time, T: Y can be chosen in the support of the packet and such that $Y^0 = T$ and $\mathrm{Im}(\psi(Y)) \neq 0$. Note that this result holds for *any* packet with support only on positive momenta – the approximately Gaussian form is not necessary.

The quantum theory of a massless scalar field in $1 + 1$ dimensional Minkowski spacetime is unphysical, not least because it suffers from an infrared divergence, so we turn our attention to $3 + 1$ dimensions. The same conclusion obtained in $1 + 1$ dimensions about persistence of the signal holds in $3 + 1$ dimensions if the wavepacket is completely spatially delocalised in the directions transverse to the direction of motion of the packet which is in the positive z-direction, say. In other words, the packet has support in momentum space on fixed values of the momentum in the x and y directions and a spread of positive values of the momentum in the z direction. In the more physical case, when the packet is localised in all three spatial dimensions and described by the state $|1_3\rangle$ in (3), the amplitude of the envelope of the packet will decay due to diffraction into the transverse directions. Thus, in $3 + 1$ dimensions the particular measure of the superluminal signal, $\mathrm{Im}(\psi(Y))$, decays as $Y^0 = T$ – the intervention time – increases. To make this concrete, let us consider the Gaussian packet $|1_3\rangle$ peaked at $\mathbf{k}_0 = (0, 0, k_0)$ and calculate $\psi(Y)$ for arbitrary points in the $z - t$ plane, $Y = (t, 0, 0, z)$. Then $\psi(Y)$ can be evaluated in closed form [2, § 3.953]:

$$\begin{aligned}\psi(Y) &= \langle 0|\hat{\phi}(t, 0, 0, z)|1_3\rangle \\ &= (\pi\sigma^2)^{-\frac{3}{4}} \int \frac{d^3k}{(2\pi)^{\frac{3}{2}}\sqrt{2|\mathbf{k}|}} e^{-\frac{(\mathbf{k}-\mathbf{k}_0)^2}{2\sigma^2}} e^{ik_\mu Y^\mu} \\ &= \frac{e^{-k_0^2/2\sigma^2}}{4\pi^{\frac{3}{4}}} \frac{e^{v_-^2/4} D_{-\frac{3}{2}}(v_-) - e^{v_+^2/4} D_{-\frac{3}{2}}(v_+)}{k_0\sigma^{-2} + iz} ,\end{aligned} \tag{6}$$

where $v_\pm = i\sigma\left[t \pm (z - ik_0/\sigma^2)\right]$ and $D_\nu(z)$ is the parabolic cylinder function.

In Figure 1, the imaginary part of $\psi(t, 0, 0, z)$ for the Gaussian 1−particle state with $\mathbf{k}_0 = (0, 0, 10)$ and $\sigma = 1$ is plotted as a function of z for $t = 0, 10, 20$. It can be seen that the amplitude of the packet envelope remains non-negligible for values of $t = Y^0$ which are a few times the spatial width of the packet. If we choose X to be at the origin, $X^\mu = (0, 0, 0, 0)$, then at each of the representative times, points in the right hand half of the support of the packet are spacelike to X. Therefore Y can be spacelike to X, in the support of the packet



FIG. 1. The imaginary part of $\psi(t, 0, 0, z)$ for the Gaussian 1−particle state with $\mathbf{k}_0 = (0, 0, 10)$ and $\sigma = 1$, plotted as a function of z for different values of $t = 0, 10, 20$.

and such that $Y^0 =$ few×(spatial extent of packet) and the measure of the superluminal signal Im($\psi(Y)$) is non-negligible. Indeed, if we let $z = t + \delta$ with $0 < \delta < 1/\sigma$, such that Y is spacelike to X but still inside the support of the wavepacket, then the asymptotic expansion of (6) for large $t \gg k_0/\sigma^2$ gives [3]:

$$\mathrm{Im}(\psi(Y)) \sim \gamma\sqrt{k_0/\sigma}\cos(k_0\delta)t^{-1} \tag{7}$$

where γ is a constant of order $\mathcal{O}(10^{-1})$. It follows that the envelope of the superluminal signal Im($\psi(Y)$) falls off like t^{-1}.

We conclude that one or more of the three interventions – the unitary kick at X, the ideal measurement of the wavepacket between $t = 0$ and $t = T$ =few×(spatial size of packet), or the measurement of $\hat{\phi}(Y)$ – is impossible. The measurement of the wavepacket is the most questionable as it is nonlocal but to begin to address these questions properly we need to model the interventions physically.

Another example of a physical quantity that cannot be measured by an ideal measurement without causing superluminal signalling is a non-abelian Wilson loop [4]. Furthermore, the observables on which the algebraic approach to relativistic quantum field theory is based are field operators smeared with test functions with bounded space*time* support and it is an open question whether or not ideal measurements of such observables would enable superluminal signalling [1].

Finally, the same no-go results apply to unitary transformations: an intervention by an external agent that produces a nonlocal unitary transformation on the state of the field, for example, a unitary transformation that rotates the state from one 1–particle state to an orthogonal 1–particle state, can enable superluminal signalling. Consider the 2−dimensional Hilbert space $\mathcal{H}$ spanned by two 1–particle states $|1\rangle$ and $|1'\rangle$. This space is a subspace of the Fock space $\mathcal{F} = \mathcal{H} \oplus \mathcal{H}^\perp$ of the scalar field and $\mathcal{H}^\perp$ denotes its orthogonal complement. The field starts in a state $|\psi\rangle = A|1\rangle + B|1'\rangle$. Consider the same 3 intervention regions, $\mathcal{O}_1$, $\mathcal{O}_2$ and $\mathcal{O}_3$, as before. A local unitary operation $e^{i\lambda\hat{\phi}(X)}$ is done at X followed by a unitary operation U done in $\mathcal{O}_2$ where

$$U = e^{i\theta}(C|1\rangle\langle 1| + D|1\rangle\langle 1'| - D^*|1'\rangle\langle 1| + C^*|1'\rangle\langle 1'|) + \mathbf{1}^\perp\,, \tag{8}$$

$|C|^2 + |D|^2 = 1$ and $\mathbf{1}^\perp$ is the identity operator on $\mathcal{H}^\perp$. Finally, $\hat{\phi}(Y)$ is measured at Y. The expected value of the measurement of $\hat{\phi}(Y)$ is

$$\langle\hat{\phi}(Y)\rangle = \langle\psi|e^{-i\lambda\hat{\phi}(X)}U^\dagger\hat{\phi}(Y)Ue^{i\lambda\hat{\phi}(X)}|\psi\rangle\,. \tag{9}$$

Differentiating w.r.t. λ and setting $\lambda = 0$ we find that the superluminal signal is nonzero if the following expression is nonzero:

$$2\mathrm{Im}\langle\psi|\hat{\phi}(X)U^\dagger\hat{\phi}(Y)U|\psi\rangle\,. \tag{10}$$

For $A = 1$, $B = 0$, $C = 0$ and $D = -1$ we find

$$\langle\psi|\hat{\phi}(X)U^\dagger\hat{\phi}(Y)U|\psi\rangle = \psi(X)^*\psi'(Y) + \psi'(X)\psi(Y)^* \tag{11}$$

where $\psi(\xi) = \langle 0\,|\hat{\phi}(\xi)|\,1\rangle$ and $\psi'(\xi) = \langle 0\,|\hat{\phi}(\xi)|\,1'\rangle$ are the one particle wave functions.

If the one particle states are single momentum states, this is easily worked out. Let us work in a box of side length L, so that

$$\hat{\phi}(X) = L^{-\frac{d}{2}}\sum_{\mathbf{k}}\frac{1}{\sqrt{2\omega_{\mathbf{k}}}}\left[a_{\mathbf{k}}u_{\mathbf{k}}(X) + a^\dagger_{\mathbf{k}}u^*_{\mathbf{k}}(X)\right] \tag{12}$$

where $u_{\mathbf{k}}(X) = e^{ik_\mu X^\mu}$. Let $|1\rangle = a^\dagger_{\mathbf{k}}|0\rangle$ and $|1'\rangle = a^\dagger_{\mathbf{k}'}|0\rangle$. Then the signal (10) evaluates to

$$\frac{L^{-d}}{\sqrt{\omega_{\mathbf{k}}\omega_{\mathbf{k}'}}}\left[\sin(k'_\mu Y^\mu - k_\mu X^\mu) - \sin(k_\mu Y^\mu - k'_\mu X^\mu)\right]. \tag{13}$$

This can be nonzero for spacelike separated X and Y. For example, for $\mathbf{k} = -\mathbf{k}'$ we obtain

$$-\frac{2L^{-d}}{\omega_{\mathbf{k}}}\sin\left[\omega_{\mathbf{k}}(Y^0 - X^0)\right]\cos\left[\mathbf{k}\cdot(\mathbf{Y} + \mathbf{X})\right], \tag{14}$$

which is nonzero for choices of spacelike separated Y and X for which $Y^0 - X^0$ is arbitrarily large. If $|1\rangle$ and $|1'\rangle$ are wave packet states, the spreading of the packets becomes relevant. However, similarly to the previous calculation, X can be chosen in the support of ψ at $t = 0$ and Y in the support of ψ' at $t = T$ and there will be superluminal signalling for T =few×(width of packet). The only unitary transformations that would definitely not violate relativistic causality are those which are products of unitaries that are perfectly localised at spacetime points on a spacelike hypersurface, e.g. $e^{i\int d^dx f(\mathbf{x})\hat{\phi}(t,\mathbf{x})}$.

III. DIFFERENT RULES

The preceding observations show that modelling external interventions on quantum fields as ideal measurements and unitary transformations in a straightforward generalisation of the text-book rules for nonrelativistic quantum theory fails. One response could be to put restrictions on the use of the formula (1) for the probabilities of outcomes of a collection of measurements.

Restricting the regions. One could require that the relation $\preceq$ between the intervention regions, $\mathcal{O}_i$, is already a partial order *before* the transitive closure is taken [1]. In the protocol considered above that would require X to be in the causal past of Y and any signalling would then be causal by fiat. This leaves the question, "What is the probability of a certain sequence of outcomes of measurement-interventions in regions which are not partially ordered?" without any answer, though there seems to be no physical reason why the relative positions of the regions in spacetime should affect agents' ability to do measurements in them. Other, even more restrictive conditions on the regions $\mathcal{O}_i$ might be considered. For example, one might require that for every pair $(\mathcal{O}_j, \mathcal{O}_k)$ such that $j < k$, either the two regions are entirely spacelike to each other or every point of $\mathcal{O}_j$ is in the causal past of every point of $\mathcal{O}_k$.[2] Once again, such rules would be silent on how to make predictions when the interventions on quantum fields take place in regions that do not satisfy these conditions.

Restricting the observables. Alternatively, one could restrict the observables, begging the question why some observables are measurable and others not. Observables which are field operators smeared with real functions over subsets of spacelike hypersurfaces (and not over time), such as those considered in [5], are essentially local and could, in principle, be measured by ideal measurements without causing superluminal signalling. Whether or not the measurements can be done in practice is another question.

Another example of an essentially local (Heisenberg) observable at time t is $a_{\mathbf{k}}(t) + a^\dagger_{\mathbf{k}}(t) = a_{\mathbf{k}} e^{-i\omega_{\mathbf{k}} t} + a^\dagger_{\mathbf{k}} e^{i\omega_{\mathbf{k}} t}$, where $\omega_{\mathbf{k}}$ is the frequency for momentum $\mathbf{k}$, which may be written as

$$\begin{aligned} & a_{\mathbf{k}}(t) + a^\dagger_{\mathbf{k}}(t) \\ & = \int d^d x\, \hat{\phi}(t,\mathbf{x}) F(\mathbf{x}) + \int d^d x\, \hat{\pi}(t,\mathbf{x}) G(\mathbf{x}) \end{aligned} \tag{15}$$

where $\hat{\pi}$ is the canonical momentum of $\hat{\phi}$, and

$$F(\mathbf{x}) = (2\omega_{\mathbf{k}})^{\frac{1}{2}} (2\pi)^{-\frac{d}{2}} \cos \mathbf{k}\cdot\mathbf{x} \tag{16}$$

$$G(\mathbf{x}) = \left(\frac{2}{\omega_{\mathbf{k}}}\right)^{\frac{1}{2}} (2\pi)^{-\frac{d}{2}} \sin \mathbf{k}\cdot\mathbf{x}\,. \tag{17}$$

To make the single momentum states normalised by working in a box of side length L, we could replace the (2π) by L in (16) and (17). Although this observable is defined on the whole constant time hypersurface at t it is a sum of truly local terms on the hypersurface and as such an ideal measurement of it will not give rise to superluminal signalling in the protocol considered above. It is "essentially local" but not "localised" and therefore unlikely to be useful for QIP in Minkowski space. Localised smearings, such as those considered in [5], have a better chance of being useful. Another example of a more localised observable is $b(t) + b^\dagger(t)$ where $b^\dagger(0)$ is defined by $b^\dagger(0)|0\rangle = |1\rangle$, a one particle wave packet state.

If

$$b^\dagger(0) = \int d^d k\, \tilde{\psi}(\mathbf{k})\, a^\dagger_{\mathbf{k}} \tag{18}$$

then the one particle wave function is

$$\psi(t,\mathbf{x}) = \int d^d k\, (2\omega_{\mathbf{k}})^{-\frac{1}{2}} (2\pi)^{-\frac{d}{2}} e^{i\mathbf{k}\cdot\mathbf{x}} e^{-i\omega_{\mathbf{k}} t} \tilde{\psi}(\mathbf{k}) \tag{19}$$

and

$$b(t) + b^\dagger(t) = \int d^d x\, \Big(\hat{\phi}(t,\mathbf{x}) J(\mathbf{x}) + \hat{\pi}(t,\mathbf{x}) K(\mathbf{x})\Big) \tag{20}$$

where

$$J(\mathbf{x}) = \int \frac{d^d k}{(2\pi)^{\frac{d}{2}}} \left(\frac{\omega_{\mathbf{k}}}{2}\right)^{\frac{1}{2}} \left[e^{i\mathbf{k}\cdot\mathbf{x}} \tilde{\psi}(\mathbf{k}) + \text{c.c.}\right] \tag{21}$$

$$K(\mathbf{x}) = -i \int \frac{d^d k}{(2\pi)^{\frac{d}{2}} (2\omega_{\mathbf{k}})^{\frac{1}{2}}} \left[e^{i\mathbf{k}\cdot\mathbf{x}} \tilde{\psi}(\mathbf{k}) - \text{c.c.}\right]\,. \tag{22}$$

$K(\mathbf{x})$ is (twice) the imaginary part of $\psi(0,\mathbf{x})$ but $J(\mathbf{x})$ is not its real part.

We see that for $b(t) + b^\dagger(t)$ to be localised, $\tilde{\psi}(\mathbf{k})$ must be such that K and J have bounded support in space. K will do so if the one-particle wavefunction ψ at $t = 0$ does but requiring J to have bounded support imposes an additional constraint and it is not clear that both constraints can be exactly satisfied. If it is sufficient for K and J to be exponentially small outside a bounded region of space then the Gaussian wave packet state (3) will do, but, strictly, to measure $b(t) + b^\dagger(t)$ in this case requires an intervention region that includes the whole spacelike hypersurface at t. If we accept that in this case intervening in a bounded region can in principle result in an ideal measurement of $b(t) + b^\dagger(t)$ to a very good approximation, it is still an open question whether it can be measured by an ideal measurement in practice.

IV. DETECTOR MODELS

A more physical approach to addressing these questions – and, for example, explaining which of the interventions considered in the first section is impossible – is to construct explicit models of the interventions and detections as interactions between the field and other quantum systems. A quantum detector for example could act as the ‘qubit’ that is manipulated by the classical agent and the quantum field coupled to it would then be part of the quantum system that performs the processing task but would not be directly intervened upon by the external agent. A detector would have a world-line or world-volume in spacetime and one would have to assume that ideal measurements and unitary transformations can be performed on it by the external agent in some region.

---

[2] This variant was described to us by Andrzej Dragan. If one wanted to interpret this rule in terms of the quantum state it seems to mean that for each $\mathcal{O}_i$ in turn, the quantum state collapses along the boundary of the future set $\{y \,|\, y \in J^+(x)\ \forall x \in \cup_{k=1}^{i} \mathcal{O}_k\}$.

One prototype of such a detector is the Unruh-Dewitt (UD) detector [6, 7]. UD detectors and their ilk couple to the field operator itself and field operators at spacelike separated positions commute. Signalling between two UD detectors will therefore not occur *if* their world-volumes are completely spacelike separated while they are switched on, and the probability for ideal measurements on the detectors themselves is given by (1). Whether UD detectors are good models of realistic detectors that can be used in QIP is an open question. In order to provide a useful qubit a UD detector needs to be switched on and off at finite times and also itself prepared and measured or manipulated at finite times. This requires the modelling of the switching process to be addressed with care [8] but at least relativistic causality can be safe with UD detectors because they couple locally to the field.

Are there other model detectors that might be applicable to QIP in Minkowski spacetime? We briefly consider two that appear in the literature. A detector that couples to one single frequency mode of a scalar field in 1+1 dimensions is investigated in [9, 10]. We can imagine that the system is in a large box of size L with periodic boundaries, say, so that the single mode state is normalisable. The Hamiltonian for the model is

$$\begin{aligned}\hat{H}(t) = w(\hat{d}^\dagger \hat{d} + \frac{1}{2}) + \Omega(\hat{f}^\dagger \hat{f} + \frac{1}{2}) \\ + \lambda(t)(\hat{d} + \hat{d}^\dagger)\left(\hat{f} e^{i\Omega x} + \hat{f}^\dagger e^{-i\Omega x}\right),\end{aligned} \tag{23}$$

where x is the position of the detector which has an internal harmonic oscillator degree of freedom with frequency w and raising operator $\hat{d}^\dagger$, and Ω is the frequency of the field mode with one-particle creation operator $\hat{f}^\dagger$. This causes superluminal signalling if interpreted as a model for a localised detector interacting with a field in Minkowski spacetime, or in a box such that the light crossing time of the box is longer than the timescale on which the model is valid.

Consider, following [10], the particular case of two static detectors at positions x_i, with frequencies w_i and couplings $\lambda_i(t)$, $i = 1, 2$. In order that the model be close to Minkowski spacetime, we assume that the box size L is large compared to $|x_1 - x_2|$. In the interaction picture, the interaction Hamiltonians are

$$\begin{aligned}H_{i,\mathrm{int}}(t) =& \lambda_i(t)(\hat{d}_i e^{-i\omega_i t} + \hat{d}_i^\dagger e^{i\omega_i t})\times \\ & \left(\hat{f} e^{i\Omega(x_i - t)} + \hat{f}^\dagger e^{-i\Omega(x_i - t)}\right)\end{aligned} \tag{24}$$

for $i = 1, 2$. The time dependence in $\lambda_i(t)$ parametrizes the switching on and off of the detectors. The interaction Hamiltonians of the two detectors at spacelike separated positions do not commute. Indeed, defining $X_i^\mu := (t_i, x_i)$ and $\Omega^\mu := (\Omega, \Omega)$, we have the commutator

$$\begin{aligned}[H_{1,\mathrm{int}}(t_1), H_{2,\mathrm{int}}(t_2)] =& -2i\lambda_1(t_1)\lambda_2(t_2)\times \\ & (\hat{d}_1 e^{-i\omega_1 t_1} + \hat{d}_1^\dagger e^{i\omega_1 t_1})\times \\ & (\hat{d}_2 e^{-i\omega_2 t_2} + \hat{d}_2^\dagger e^{i\omega_2 t_2})\times \\ & \sin\Omega_\mu (X_1 - X_2)^\mu\,.\end{aligned} \tag{25}$$

This is nonzero for almost every pair of spacetime points along the trajectories of two detectors which are spacelike separated. This means that the unitary evolution operator in the interaction picture does not factorise into a product of an evolution operator for detector 1 and one for detector 2 and this leads to superluminal signalling between the detectors as we show explicitly below.

Let both detectors and the scalar field be initially in their free ground states at $t = 0$ and assume that the detectors are switched off before $t = 0$: $\lambda_i(t) = 0$ for $t < 0$. Working in the interaction picture, at time $T > 0$ we measure for detector 1 the expectation value, denoted $E_1(T)$, of the operator

$$\mathcal{O}(T) = \omega_1(\hat{d}_1(T)^\dagger \hat{d}_1(T) + \frac{1}{2}) = \omega_1(\hat{d}_1^\dagger \hat{d}_1 + \frac{1}{2}). \tag{26}$$

$E_1(T)$ ought to be independent of λ_2 so long as detector 1 remains outside the causal future of the point $(0, x_2)$, i.e. when $T < |x_2 - x_1|$. Otherwise, an observer at (T, x_1) could use their detector to infer whether spacelike separated detector 2 has been switched on or not (i.e. whether $\lambda_2 \neq 0$ or not). We do not carefully model the switching process and simply take $\lambda_i(t)$ to be a step function, constant and nonzero only in the interval $0 < t \leq T$ and then $E_1(T)$ can be evaluated explicitly using the Heisenberg Picture analysis presented in [10]. It can be shown that $E_1(T)$ does in general depend on λ_2, and by way of example let us evaluate it for a specific set of values: $w_1 = w_2 = \Omega = 1$, $\lambda_1 = \frac{1}{2}$, $x_1 = 0$ and $x_2 = 2\pi$. With a constant coupling $\lambda_2(t) = \lambda_2$ for $0 < t \leq T = \sqrt{2}\pi < |x_2 - x_1|$ one obtains

$$E_1(\sqrt{2}\pi) = \frac{1}{4}\left(2 + \pi^2\right) + \frac{1}{3}\pi^4 \lambda_2^2 + \mathcal{O}(\lambda_2^4). \tag{27}$$

We see a second–order dependence on the coupling constant λ_2 of detector 2 – a superluminal signal.

This signalling becomes more comprehensible if the detector is interpreted as a variant of an UD detector in which the detector couples both to the field and to its conjugate momentum. Indeed, modifying the calculation that leads to (15), we find for the factor in the interaction hamiltonian $H_{1,\mathrm{int}}(t)$ for the first detector

$$\begin{aligned}& \hat{f} e^{i\Omega(x_1 - t)} + \hat{f}^\dagger e^{-i\Omega(x_1 - t)} \\ & \quad = \int dx \hat{\phi}(t, x) F_1(x) + \int dx \hat{\pi}(t, x) G_1(x)\end{aligned} \tag{28}$$

where

$$F_1(x) = (2\Omega)^{\frac{1}{2}} L^{-\frac{1}{2}} \cos[\Omega(x - x_1)] \tag{29}$$

$$G_1(x) = \left(\frac{2}{\Omega}\right)^{\frac{1}{2}} L^{-\frac{1}{2}} \sin[\Omega(x - x_1)]\,. \tag{30}$$

We can see that the effective spatial extension of the detector is not bounded: the detector is not localised and two detectors which are nominally spacelike separated actually overlap over all space.

In [11] a more ambitious model of a detector defined using additional quantum fields coupled to the field being measured is considered. Since the interaction between the fields is local no superluminal signalling can be enabled as a result of the interaction in and of itself. If taken literally, the effective model of the detector given in equation (14) in [11] looks like it does enable superluminal signalling since the interaction Hamiltonians of two detectors centred at spacelike separated positions do not commute. Just as in the calculation above, this will result in the response of one detector depending on whether or not there exists a second one spacelike to it. Again, this happens because detectors constructed as described in [11] cannot be genuinely spacelike separated: they may be centred around localised positions but will always have some spatial overlap. The authors themselves state that the detector is nonlocal. However, the question of whether the state of such a detector itself is measurable by an ideal measurement immediately arises: does an ideal measurement on the non-local detector enable superluminal signalling? This is not addressed in [11] but it should not be hard to determine.

V. CAVITIES

Those familiar with cavity QED might be surprised at these results. After all, in a cavity, ideal measurements of observables such as particle number *can* be done [12] and there's a well-known, successful model of an atom-qubit interacting with QED in a cavity which is of the form investigated above in which a detector couples to a single mode of the field: the Jaynes-Cummings model (see *e.g* [13] for a review). The reason there is no conflict with the results presented here is that the Jaynes-Cummings model is a phenomenological model which applies only on time scales many orders of magnitude larger than the light crossing time of the cavity. It does not describe the physics on time scales of order the light crossing time or shorter.[3] If there are two atoms in a cavity, they cannot be placed further apart than the size of the cavity and so their worldlines cannot remain spacelike to each other on the time scales on which the Jaynes-Cummings model applies and so no possibility of superluminal signalling arises in the model.

The Jaynes-Cummings Hamiltonian and its relatives cannot model an atom-qubit coupled to a quantum field in Minkowski spacetime, or in any spacetime where two atom-qubits can be placed at distances larger than the timescale on which the detector model is valid. A model detector such as that considered in [9, 10] would not do violence to relativistic causality if it were interpreted phenomenologically and confined to a spacetime of much smaller spatial extent than the timescale on which the model is valid since this would effectively impose a non-relativistic causal order on the intervention-events in the cavity. In the light of the phenomenological successes of such models, one might be tempted to declare more widely that QFT is only physical in a box and that intervention regions $\mathcal{O}_i$ must always have extent in time at least the light-crossing time of the box, imposing a non-relativistic causal order on interventions on the field. Such a rule would be highly restrictive and could not accommodate the questions that arise when the operations of the external agents on a quantum field take place in arbitrary, quasi-local spacetime regions, exactly the sort of situation that a genuinely relativistic approach to QIP aims to describe.

VI. CONCLUDING REMARKS

The questions remain. What are accurate models of physical interventions on relativistic quantum fields in quasi-local regions of Minkowski spacetime and what applications might they have in QIP? Localised UD-type detectors form one class of models: Are they realistic? Are there others? The questions are interesting from the point of view of QIP but also because the attempt to answer them pushes us to address more foundational issues. The struggle to describe measurement of relativistic quantum fields in a physical way reveals the limitations of the canonical and operational approach to quantum theory. Even when a detector is modelled as a quantum system, to be a useful qubit one still has to rely on the assumption that an external agent can measure *it* by an ideal measurement and/or do unitary transformations on it. But a more fundamental description of the detector – and indeed the agent – would be in terms of quantum field theory and so one is not solving the problem but just pushing it one step away: how would one model the effect of an ideal measurement on the detector if the detector were described within a quantum field theory (*cf.* [11])?

A more physical approach is needed: we require a framework for closed relativistic quantum systems including detectors, in which experimental, measurement-like situations can be analysed fully quantum mechanically in an essentially relativistic way.

---

[3] The laboratory values of the Rabi frequency and cavity size given in [13] mean that the timescale for the model is about 10^5 times the light crossing time of the cavity. Any result based on the assumption that the dynamical evolution of the state of an atom in a cavity is described by a Jaynes-Cummings type model on time scales much shorter than the Rabi time period, such as [14] for example, is unphysical.

One proposal for such a framework takes the relativistic branch of the 'fork in the road' set out by Dirac early in the history of quantum mechanics. In the 1933 paper, "The Lagrangian in quantum mechanics," Dirac wrote that the Lagrangian approach to classical mechanics is probably more fundamental than the Hamiltonian approach because the former is relativistically invariant whereas the latter is "essentially nonrelativistic" [15]. In quantum theory the Hamiltonian approach leads to canonical quantisation, Hilbert space, operators, observables, transformation theory and the textbook rules for predicting the outcomes of measurements of observables and for the collapse of the state after an ideal measurement. These aspects of the canonical theory are indeed more or less divorced from the spacetime nature of the physical world revealed by relativity.

The relativistic alternative is to base quantum mechanics on the Lagrangian approach to classical mechanics and Dirac showed that this leads to the path integral[15]. The path integral roots quantum theory firmly in spacetime – rather than Hilbert space – as the arena for physics. In a path integral or sum-over-histories approach, the physical world is described directly in terms of events in spacetime and Feynman's famous paper on the path integral is aptly titled "The spacetime approach to non-relativistic quantum mechanics"[16]. The path integral approach to the foundations of quantum theory has been championed in more recent times by Hartle [17, 18] and by Sorkin [19]. In [20] measurement situations in closed quantum systems are analysed from a sum-over-histories perspective, with an emphasis on a condition of decoherence. In [21] a solution of the "measurement problem" of quantum mechanics is set out which is valid, at least, when there are certain sorts of permanent records of measurement outcomes. As a framework for closed quantum systems which deals directly with spacetime events, the path integral approach is eminently suitable for the investigation of measurements on relativistic quantum fields in Minkowski spacetime.

ACKNOWLEDGMENTS

We thank Andrzej Dragan, Jorma Louko and Alejandro Satz for useful and stimulating discussions. Research at Perimeter Institute for Theoretical Physics is supported in part by the Government of Canada through NSERC and by the Province of Ontario through MRI. This research was supported in part by the National Science Foundation under Grant No. NSF PHY11-25915 and by COST Action MP1006. DMTB is supported by an EPSRC Studentship. The work of LB is supported by an Imperial College Junior Research Fellowship.

---

[1] R. D. Sorkin, in *Directions in General Relativity: Proceedings of the 1993 International Symposium, Maryland, Vol. 2: Papers in honor of Dieter Brill*, edited by B. Hu and T. Jacobson (CUP, Cambridge, 1993), pp. 293–305, arXiv:gr-qc/9302018.

[2] I. Gradshteyn, I. Ryzhik, A. Jeffrey, and D. Zwillinger, *Table of Integrals, Series, And Products* (Academic Press, 2007), ISBN 9780123736376, `http://www.mathtable.com/gr/`.

[3] *NIST Digital Library of Mathematical Functions*, `http://dlmf.nist.gov/12.9.1`, `http://dlmf.nist.gov/12.9.3`, accessed: 21/06/2012.

[4] D. Beckman, D. Gottesman, A. Kitaev, and J. Preskill, Phys. Rev. **D65**, 065022 (2002), arXiv:hep-th/0110205.

[5] M. Zych, F. Costa, J. Kofler, and C. Brukner, Phys.Rev. **D81**, 125019 (2010), arXiv:1003.3354.

[6] W. G. Unruh, Phys. Rev. **D14**, 870 (1976).

[7] B. S. DeWitt, in *General Relativity; an Einstein centenary survey*, edited by S. Hawking and W. Israel (Cambridge University Press, 1979).

[8] J. Louko and A. Satz, Class.Quant.Grav. **25**, 055012 (2008), arXiv:0710.5671.

[9] E. Martin-Martinez, I. Fuentes, and R. B. Mann, Phys.Rev.Lett. **107**, 131301 (2011), arXiv:1012.2208.

[10] A. Dragan and I. Fuentes (2011), arXiv:1105.1192.

[11] F. Costa and F. Piazza, New J.Phys. **11**, 113006 (2009), arXiv:0805.0806.

[12] B. R. Johnson, M. D. Reed, A. A. Houck, D. I. Schuster, L. S. Bishop, E. Ginossar, J. M. Gambetta, L. DiCarlo, L. Frunzio, S. M. Girvin, et al., Nat. Phys. **6**, 663 (2010), arXiv:1003.27345.

[13] R. Miller, T. E. Northup, K. M. Birnbaum, A. Boca, A. D. Boozer, and H. J. Kimble, Journal of Physics B: Atomic, Molecular and Optical Physics **38**, S551 (2005).

[14] E. Martin-Martinez, A. Dragan, R. Mann, and I. Fuentes (2011), arXiv:1112.3530.

[15] P. A. M. Dirac, Physikalische Zeitschrift der Sowjetunion **3**, 64 (1933).

[16] R. P. Feynman, Rev. Mod. Phys. **20**, 367 (1948).

[17] J. B. Hartle, Vistas Astron. **37**, 569 (1993), arXiv:gr-qc/9210004.

[18] J. B. Hartle, in *Proceedings of the Les Houches Summer School on Gravitation and Quantizations, Les Houches, France, 6 Jul - 1 Aug 1992*, edited by J. Zinn-Justin and B. Julia (North-Holland, 1995), arXiv:gr-qc/9304006.

[19] R. D. Sorkin, Mod. Phys. Lett. **A9**, 3119 (1994), arXiv:gr-qc/9401003.

[20] J. B. Hartle, in *Directions in General Relativity: Proceedings of the 1993 International Symposium, Maryland, Vol. 2: Papers in honor of Dieter Brill*, edited by B. Hu and T. Jacobson (CUP, Cambridge, 1993), arXiv:gr-qc/9301011.

[21] R. D. Sorkin, in *Foundations of Space and Time*, edited by G. Ellis, J. Murugan, and A. Weltman (Cambridge University Press, 2010), arXiv:1004.1226.